

Degree of mode localization in Random lasing from ZnO nanoparticles

**Johannes Fallert, Janos Sartor, Roman J. B. Dietz,
Daniel Schneider, Victor Zalamai, Claus Klingshirn, Heinz Kalt**

<https://doi.org/10.1364/CLEO.2009.CTuW2>

Abstract

We present an experimental procedure to directly extract the degree of localization of random laser modes in ZnO powders. Strongly localized and extended modes are found to coexist within the same ensemble area.

References:

1. D. S. Wiersma, P. Bartolini, A. Lagendijk, and R. Righini, "Localization of light in a disordered media," *Nature* 390, 671 (1997).
2. H. Cao, Y. G. Zhao, S. T. Ho, E. W. Seelig, Q. H. Wang, and R. P. H. Chang, "Random laser action in semiconductor powder," *Phys. Rev. Lett.* 82, 2278 (1999).
3. D. S. Wiersma, "The physics and applications of random lasers," *Nature Physics* 4, 359 (2008).
4. C. Vanneste, and P. Sebbah, "Selective excitation of localized modes in active random media," *Phys. Rev. Lett.* 87, 183903 (2001).
5. C. Vanneste, P. Sebbah, and H. Cao, "Lasing with resonant feedback in weakly scattering random systems," *Phys. Rev. Lett.* 98, 143902 (2007).