

## **PROSPECTS FOR THE UTILIZATION OF THE *Silphium perfoliatum* AND *Silphium integrifolium* FOR RENEWABLE ENERGY PRODUCTION IN MOLDOVA**

DOROFTEI VEACESLAV<sup>1</sup>, ȚÎȚEI VICTOR<sup>1</sup>, ABABII  
ALEXEI<sup>1</sup>, ANDREOIU ANDREEA CRISTINA<sup>2</sup>, COZARI  
SERGHEI<sup>1</sup>, GADIBADI MIHAI<sup>1</sup>, CEREMPEI VALERIAN<sup>1</sup>,  
GUDIMA ANDREI<sup>3</sup>, NAZAR BORIS<sup>3</sup>, DARADUDA  
NICOLAE<sup>3</sup>

<sup>1</sup>“Alexandru Ciubotaru” National Botanical Garden (Institute),  
Chișinău, Republic of Moldova, ababiialexei@gmail.com,  
biocom04@gmail.com, vic.titei@gmail.com,  
csv2007@yahoo.com, mm.gadibadi@gmail.com

<sup>2</sup>Research - Development Institute for Grassland Brașov,  
Romania, andreea.andreoiu@pajisti-grassland.ro

<sup>3</sup>Technical University of Moldova, Chișinău, Republic of  
Moldova, andrei.gudima@tran.utm.md,  
nicolae.daraduda@tran.utm.md

The cup plant *Silphium perfoliatum* L. and rosinweed *Silphium integrifolium* Michx., *Asteraceae* family, native to North America, which were introduced and cultivated in the experimental plot of the “Alexandru Ciubotaru” National Botanical Garden (Institute) of Moldova served as subjects of the research. This research was aimed at evaluating the biomass energy indices of green harvested mass as feedstock for biogas production and stem dry mass as feedstock for cellulosic ethanol. The biochemical composition of green mass and dry stem mass have been determined by NIRS technique PERTEN DA 7200 of the Research and Development Institute for Grassland Brașov, Romania. It has been determined that harvested green mass from *Silphium* species contained: 13.7-16.4 % CP, 84-90 g/kg ash, 48-49 g/kg ADL, 306-327 g/kg Cel, 222-227 g/kg HC and the estimated biochemical methane potential of studied

substrates were 331-338 l/kg ODM. The analysis of cell wall components of stem dry mass revealed that studied substrates from *Silphium* species contained 487-534 g/kg cellulose, 258-274 g/kg hemicellulose, 95-112 g/kg acid detergent lignin and the estimated theoretical ethanol yield averaged 553-575 L/t. The investigated *Silphium integrifolium* and *Silphium perfoliatum* species, may be use as multi-purpose feedstock for renewable energy production in Republic of Moldova.

Financial support for research was offered by the National Agency for Research and Development, project no. 20.80009.5107.02

**Key words:** biochemical composition, biomethane potential, dry biomass, green mass, *Silphium integrifolium*, *Silphium perfoliatum* theoretical ethanol potential